REVIEW ARTICLE

CHARACTERISTICS, DETECTION AND TYPING METHODS OF COMMUNITY ACQUIRED METHICILLIN RESISTANT STAPHYLOCOCCUS AUREUS (CA-MRSA) - A REVIEW

Lia Natasha Amit¹, Fong Siew Moy² and Daisy Vanitha John¹

¹Biotechnology Research Institute, Universiti Malaysia Sabah, Jalan UMS, 88400, Kota Kinabalu, Sabah, Malaysia

²Infection Control Unit, Hospital Wanita dan Kanak-Kanak Sabah, Karung Berkunci 187, 88996, Kota Kinabalu, Sabah, Malaysia

Corresponding author:

Daisy Vanitha John

Email: daisyvanitha@gmail.com

ABSTRACT

Staphylococcus aureus are gram positive cocci which colonize the skin and mucous membranes particularly the anterior nares. Prevalence of nosocomial infections associated with methicillin resistant *S. aureus* have been reported in hospitals (HA-MRSA) for over five decades. Recently, community-acquired MRSA (CA-MRSA) has emerged as a cause of skin and soft tissue infections in healthy individuals. These strains are sensitive to antimicrobials, carry genes for Panton-Valentine leukocidin (PVL) toxin and belong to the staphylococcal cassette chromosome (SCC) *mec* type IV or V. The suspected mode of transmission involves close contact with carriers leading to skin or nasal colonization that result in subsequent active infection. Molecular typing is used to determine the mode of transmission of CA-MRSA in the community. General typing methods such as pulsed-field gel electrophoresis (PFGE) and multilocus sequence typing (MLST) and specific methods for *Staphylococci* such as SCCmec typing and *spa* typing have the capability to characterize bacterial chromosomes and mobile genetic elements. Combination of these molecular typing methods is necessary as each method has its own advantages with respect to discriminatory power, rapidity, cost effectiveness, reproducibility, and ease of performance.

Key words: CA-MRSA, prevalence, transmission, PVL toxin, SCCmec, *spa* typing, PFGE, MLST

INTRODUCTION

Staphylococcus aureus is a causative agent of both suppurative diseases such as folliculitis, furuncle, carbuncle, impetigo and metastatic infections such as osteomyelitis, septic arthritis, and pneumonia. *S. aureus* possesses toxins which are responsible for food poisoning and toxic shock syndrome. Bacteremia and endocarditis are examples of staphylococcal infections that can lead to fatality¹. Penicillin was effective to treat staphylococcal infections until penicillin-resistant *S. aureus* were detected². Methicillin was introduced as a semisynthetic drug to ward off the penicillin resistant strains. Methicillin-Resistant *S. aureus* (MRSA) strains emerged after two years of methicillin usage and they were primarily associated with nosocomial infections and termed as hospital acquired-MRSA (HA-MRSA)^{3,4}. In the early 1990s, MRSA infections indicated an altered trend in which individuals without healthcare associated risk factors developed infections. These infections are believed to be acquired in the community and caused by strains referred as Community-acquired MRSA (CA-MRSA).

Definition of CA-MRSA

The definition of CA-MRSA infection has been proposed by Centers for Disease Control and Prevention (CDC). A CA-MRSA must be isolated from an outpatient or isolated within 48-72 hours of hospital admission. In addition, the patient must not have the history of previous hospitalization, antimicrobial therapy within past 6 months, surgery and dialysis, medical devices or catheters that are permanently attached through the patient's skin, and family members exposed to healthcare settings⁵⁻⁸.

Prevalence of CA-MRSA:

The first CA-MRSA cases were reported from the indigenous Aboriginal population from Western Australia⁹. In 1997, CDC reported a CA-MRSA outbreak among four pediatric patients in Minnesota and North Dakota, USA, all suffered from pneumonia. The MRSA isolates obtained from all these patients were susceptible to antibiotics other than beta-lactams¹⁰. CA-MRSA infection also reported among native American villagers and the risk factor of infection was attributed to sharing contaminated saunas and antibiotic treatment¹¹. High prevalence of CA-MRSA skin and soft tissue

infections among Pacific Islanders was also reported in Hawaii¹².

The prevalence of MRSA in the community and the hospital settings was studied in eight Asian countries from September 2004 to August 2006 that participated in ANSORP, the Asian Network for Surveillance of Resistant Pathogens. The overall HA-MRSA and CA-MRSA rates were 67.4% and 25.5% respectively. The highest prevalence of CA-MRSA was reported in Sri Lanka (38.8%), Taiwan (34.8%), Vietnam (30.1%) and Philippines (30.1%). The CA-MRSA prevalence in Korea, Hong Kong, India, and Thailand were 15.6%, 8.5%, 4.3%, and 2.5% respectively. The CA-MRSA were predominantly isolated from pus samples (70.5%) and less likely from blood (9.1%), sputum (8.6%), and urine (4.3%)^{13,14}. The first CA-MRSA case in Malaysia was documented in 2008 and the strains isolated since then were reported to carry SCCmec IV or V and Panton-Valentine leukocidin (PVL) toxin genes¹⁵⁻¹⁸. The preliminary alert organism surveillance data of Hospital Wanita Dan Kanak-Kanak Sabah (HWKKS), Sabah, Malaysia showed 35 (70%) and 49 (86%) CA-MRSA cases in 2014 and 2015 respectively of the total MRSA cases showing an increasing trend in recent years (unpublished data). The exact route of transmission has not been identified in the above cases. However, close contact with carriers leading to skin or nasal colonization and subsequent active infection might be considered as the source of infection.

CA-MRSA associated infections

Younger patients with no healthcare associated risk factors are more likely to suffer from CA-MRSA infections. The common clinical symptoms of the CA-MRSA among children were abscess, cellulitis, and pneumonia^{6,19}. The incidences of pyomyositis and myositis cases caused by CA-MRSA strain, USA300 have been shown to have increased in a 5 years period in Texas Children's Hospital²⁰. CA-MRSA infections were also shown to be associated with Waterhouse-Friderichsen syndrome, necrotizing fasciitis, purpura fulminans etc²¹⁻²³.

Characteristics of CA-MRSA isolates

CA-MRSA strains are found to be smaller in size compared to HA-MRSA due to their less metabolic burden on protein synthesis during replication. They have a faster growth rate which facilitates successful colonization in humans. The CA-MRSA isolates were shown to be more susceptible to antimicrobials compared to nosocomial isolates²⁴. Molecular analysis of CA-MRSA isolates from different regions including USA, France, Switzerland, New Zealand, Australia, and Western Samoa revealed that they share same genetic

characteristics, such as harboring SCCmecIV or V element and PVL toxin²⁵. It is believed that *S. aureus* strain was initially infected by a phage harboring PVL genes (lukS-PV and lukF-PV) and integrated into the chromosome and became a prophage. Later, horizontal transfer and insertion of SCCmec element which consists of *mecA* gene into the chromosome led to the existence of CA-MRSA²⁶.

1.SCCmec element:Staphylococcal cassette chromosome (SCC) elements are genomic islands that serve as machineries to capture foreign DNA segments in order to survive in harsh environments. SCCmec element was recognized after two recombinase genes, *ccrA* and *ccrB*, encoding recombinase A and B respectively were identified while nucleotide sequencing of the entire *mecDNA* of MRSA strain, N315. The recombinase A and B are responsible for the excision and integration of SCCmec element into *S. aureus* chromosome^{27,28}. SCCmec element encodes methicillin-resistance which is a modified penicillin-binding protein. Based on International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements (IWG-SCC), SCCmec element consists of *ccr* gene complex and *mec* gene complex. The SCCmec types are determined by a combination of *ccr* gene complex and *mec* gene complex. The *ccr* gene complex consists of *ccr* genes and surrounding open reading frames with unknown function. The *mec* gene complex consists of *mecA* gene, *mecI* and *mecR1* gene (regulatory genes), and 20 to 45 kb of *mec*-associated DNA (insertion sequences). SCCmec typing shows that the CA-MRSA strains generally carry SCCmecIV or V elements where SCCmecIV possesses Type 2 *ccr* gene and Class B *mec* gene complex while SCCmecV possesses Type 5 *ccr* gene and Class C *mec* gene complex^{24,29}. The SCCmecIV element is further divided into several subtypes based on the differences in J1 region found between the right chromosomal junction and the *ccr* complex³⁰.

2.Panton-Valentine leukocidin (PVL) toxin: Panton-Valentine leukocidin (PVL) is a cytotoxin that causes leukocyte destruction and tissue necrosis and has become one of the constant markers for CA-MRSA. PVL toxin is composed of two classes of proteins, class F (LukF-PV) and class S (LukS-PV), they bind together into a heptamer and form pores on the polymorphonuclear leukocytes membranes²⁶.

Previous findings show that *S. aureus* isolates that carry PVL toxin genes were linked with furunculosis, community-associated pneumonia, acute hematogenous osteomyelitis, myositis and pyomyositis particularly in younger patients^{31,32}. Occurrence of sepsis and coagulopathy caused by

PVL positive CA-MRSA strains were reported among adolescents^{33,34}. Patients who were infected with the PVL positive CA-MRSA strains required more surgical drainage procedures and needed more attention in ICU²⁰. Pneumonia associated with the PVL positive *S. aureus* was considered lethal because of the lower survival rate after hospitalization³².

***Staphylococcus aureus* colonizes particularly anterior nares**

S. aureus is the normal microflora of the human skin and colonizes particularly the nose, axillae, perineum, back of the wrist and toeweb³⁵⁻³⁷. The colonization rates were reported to be 9.2%, 9.9%, and 20.7% in vagina, labia minora, and anterior nares respectively showing that the anterior nares were more likely to be colonized with a higher percentage of this bacterium³⁸. The presence of large apocrine glands in anterior nares and perineum gives a suitable environment for staphylococci to live³⁹. Moreover, nasal vestibule (vestibule nasi) or anterior nares are lined with keratinized, stratified squamous epithelium, sweat glands, and sebaceous glands and free from the host defenses thus serves as a suitable spot for staphylococci to reside. Nasal carriage rate was shown to be higher when the nasal swab samples were obtained from the nasal vestibule compared with the mucous membrane of the nasal fossa⁴⁰.

Colonization *S. aureus* is divided into 3 types; persistent carrier, intermittent carrier, and noncarrier. Persistent carriers consist of 20% in the general population and they normally carry only one type of strain. They might gain the same staphylococcus strain from the other body sites or from the family contacts⁴¹. Sixty percent of the population comprised of intermittent carriers where a certain type of strain colonizes them temporarily before replaced by another type of strain. Only 20% population is categorized as noncarriers⁴². CA-MRSA colonization has been reported among healthy children attending day care centers, kindergartens and pre-schools in Korea, Hong Kong and Taiwan⁴³⁻⁴⁵.

Bacterial and host factors influence *Staphylococcus* colonization

The interaction between the host cell membrane and the bacterial cell wall component is essential for successful colonization which is influenced by several factors. Bacterial extracellular structures facilitate the bacterial adherence to the nasal mucosal cells. Fibronectin-binding protein of *S. aureus* functions by binding to the human fibronectin while teichoic acid was shown to inhibit the adherence of staphylococci to the host

epithelial cell^{46,47}. Bacterial interference by decolonization of *S. aureus* in newborns shown to significantly increase colonization by gram-negative bacilli⁴⁸.

S. aureus shows high affinity and binding to the nasal mucosal cells of persistent carriers as compared to noncarriers⁴⁹. Host factors such as age, race, histocompatibility antigens (HLA), nasal abnormalities, bacterial/viral infections and sex hormones were shown to be associated with *S. aureus* colonization. Children are more inclined to be persistent carriers compared to adults. The persistence of *S. aureus* carriage was also observed among people between 10 to 20 years old. Adults are generally intermittent carriers of *S. aureus*⁵⁰. Caucasians have a higher rate of *S. aureus* nasal colonization compared to black Africans⁵¹. Individuals with HLA-DR3 type was linked with *S. aureus* nasal carriage, while individuals with HLA-DR2, HLA-DRI and HLA-Bw35 were less likely to be carriers⁵². Healthy people with nasal abnormalities such as damaged turbinate and deviated septum display high rates of *S. aureus* nasal colonization⁴⁰. People who were affected by a viral infection in the upper respiratory tract display greater adherence of *S. aureus* due to alteration of the nasal mucosal cells⁵³. Previous hospitalization, previous antimicrobial treatment with cotrimoxazole, and lower CD4 counts in HIV patients are shown to be the risk factors of MRSA colonization among tuberculosis patients⁵⁴. Lastly, women with high karyopyknotic index (KI) or estrogen level are prone to *S. aureus* nasal carriage⁵⁵.

Transmission of CA-MRSA:

Familial transmission occurs when two or more members of a family who live in the same postal address are colonized with MRSA strains that have similar genotype. CA-MRSA disseminates from children to parents or vice versa. Transmission is also possible between families as it has been shown that two families who lived in the same neighbourhood were found to be colonized with similar CA-MRSA strains⁵⁶⁻⁵⁸. Dissemination of CA-MRSA strain in the community setting encircling 26 households was reported in Denmark. Dissemination of CA-MRSA also occurs from the workplace to the household or vice versa. CA-MRSA strains might also disseminate from the other countries⁵⁹. The following five risk factors for CA-MRSA (5 C's) transmission are proposed by CDC. They are cleanliness (poor hygiene), compromised skin integrity, frequent skin contact (close contact), contaminated surfaces or shared items, and crowding or frequent antibiotic usage or overuse of antibiotics⁶⁰.

The multidrug resistant USA300 clone in homosexual men was reported in San Francisco causing skin and

soft tissue infection (SSTI), including folliculitis, cellulitis, and abscesses⁶¹. CA-MRSA skin and soft tissue infection was reported in a college football team and the risk factors were shown to be sharing towels, bar soaps and delaying the treatment of cuts and abrasions⁶². CA-MRSA cases were reported among prisoners in USA and the risk factors involved were long period of stay, history of antibiotic use, crowding, unhygienic condition, and comorbidities. The military population also shared similar risk factors of CA-MRSA infections and the additional risk factors included were sharing crowded barracks and physical trauma associated with the recruitment training⁶³.

CA-MRSA infections were reported in two child centers involving an index patient with pneumonia and preseptal cellulitis and colonization among other children who were MRSA asymptomatic carriers which was confirmed by similar PFGE pattern. The risk factor of the MRSA colonization was due to two to three courses of oral antibiotic consumption and closed contact between children⁶⁴.

CA-MRSA has started to spread from the community to hospitals, neonatal intensive care units (NICUs), nurseries, maternity settings and among healthcare workers^{13,65-68}. CA-MRSA carriage among newborns without risk factors was reported in the UK healthcare setting⁶⁹. A clinical case of CA-MRSA transmission from father to infant was reported in the intensive care unit in Saudi Arabia⁷⁰. CA-MRSA colonization was also observed among preclinical and clinical students⁶⁸. Predisposing factors for *S. aureus* colonization were shown to be smoking, chronic sinusitis, usage of antibiotics in the past 3 months, and utilization of the hospital entrance^{45,71,72}.

Methods for detecting CA-MRSA transmission

Screening or active surveillance of CA-MRSA is conducted by collecting the nasal swab sample from an individual who is not presenting with any symptoms of disease. The sample is then processed in the laboratory to determine whether the individual is a MRSA asymptomatic carrier⁷³. The implementation of rapid diagnostic methods with high sensitivity is crucial to detect the colonization⁷⁴. Typing is necessary for epidemiological investigations to determine the source and mode of transmission of CA-MRSA strains in the community⁷⁵. Currently, there is no single ideal typing method available and a combination of different typing methods has been used to understand the transmission of MRSA in the community^{76,77}.

Conventional methods: Culture and antibiotic susceptibility testing are used for the identification of CA-MRSA. Introduction of chromogenic agar excludes the requirement of subculture from the primary isolation plate and biochemical tests for MRSA identification thus help to minimize the time. The chromogenic agar such as Chromogenic MRSA/Denim Blue agar, oxacillin resistance screening agar base, MRSA Select CHROM agar, MRSA Ident agar are more expensive than the media used in conventional microbiological methods⁷⁸. The antibiotic susceptibility tests show that the MRSA isolates identified from the patients without risk factors were susceptible to erythromycin, clindamycin, gentamicin, and trimethoprim-sulfamethoxazole⁶. Rapid detection of ciprofloxacin-resistant MRSA is possible within five hours by using BacLite *Rapid MRSA* (Acolyte Biomedica, UK) where ATP production is quantified by bioluminescence⁷⁹. The above phenotypic methods have limited discriminatory power and poor typeability.

***mecA* gene PCR:** Detection of MRSA colonization among patients was greatly improved when PCR is performed for *mecA* gene⁸⁰. The time allocated for the detection is reduced from four days to one day. The qMRSA (multiplexed-real time PCR) detects *mecA* and *femA* genes from *S. aureus*, and only *femA* gene from *S. epidermidis* and thus the sample which contained MRSA and *S. epidermidis* can be easily distinguished⁸¹.

Molecular typing methods

Genetic knowledge has been applied in the clinical management in order to determine the source of infection. Molecular characteristics of the bacterial isolates found in patients and healthy individuals, especially the family contacts are compared. If the molecular characteristics of the bacterial isolates are similar, then the transmission of the single strain was found to have occurred⁸². Various molecular typing methods have been used such as pulse-field gel electrophoresis (PFGE), multi locus sequence typing (MLST), SCC*mec* typing and spa typing^{83,84}.

Pulse field gel electrophoresis (PFGE): PFGE is considered as the 'gold standard' among molecular typing methods for *S. aureus* due to its excellent discriminatory power and intra-laboratory reproducibility. Restriction endonucleases spot the specific sites on the bacterial chromosome and digest the bacterial DNA into large DNA fragments. These large DNA fragments are then separated by gel electrophoresis by electric current which is periodically switched in different directions. If the bacterial isolates display similar band patterns in the gel, then the bacterial isolates are considered genetically related^{76,77,85}. PFGE clearly

discriminated MRSA and methicillin susceptible *S. aureus* (MSSA) isolates that differed by just two bands in pediatric patients with pneumonia⁸⁶. The method is technically demanding, labour-intensive, time-consuming, with limited interlaboratory portability and it may lack the resolution power to distinguish bands of nearly identical size^{76, 87}.

Multilocus sequence typing (MLST): MLST involves PCR amplification and sequencing of seven housekeeping genes (*arcC*, *aroE*, *glpF*, *gmk*, *pta*, *tpi*, and *yqiL*). The sequences at each of the seven *loci* are compared with all the known alleles at that *locus*, and a number representing a previously described allele or a new one is assigned to the *locus*. For a given isolate, alleles present at each gene position are combined into an allelic profile and assigned a sequence type (ST) designation⁷⁷. Closely related isolates have identical STs, or STs that differ at a few *loci*, whereas unrelated isolates have unrelated STs. PCR is simple and cheap and MLST data has interlaboratory portability with standard nomenclature. However, the MLST has only moderate discriminatory power, too labor-intensive, and time consuming^{24,76,87}.

SCCmec typing by multiplex PCR: SCCmec typing of the MRSA isolates is based on the combination of *mec* gene complex and *ccr* gene complex. PCR amplification is performed with specific primers which target the oligonucleotide sequences of the SCCmec elements⁸⁸. SCCmec typing has been improved by the introduction of novel multiplex PCR primers that are specific for SCCmec subtypes which become cost and time-efficient⁸⁹. The methods are relatively straight forward, simple to perform, and inexpensive. However, it can be less useful in investigating local outbreaks, as the SCCmec elements are generally stable⁷⁷.

Staphylococcus aureus protein A (spa) typing: Protein A is the virulent factor in *S. aureus* which binds nonspecifically to the Fc region of the antibody. The hypervariable X region of *spa* gene which encodes for protein A become one of the steadfast molecular typing methods in certain countries such as Belgium and Germany for MRSA active surveillance because of its rapidity and high throughput^{90,91}. Initially, the X region of the *spa* gene consists of different short sequence repeat regions (SSRRs) is amplified by PCR followed by sequencing. Each SSRR contains about 24bp and assigned an "r" number. The software for *spa* type, Ridom StaphType is used to classify and determine the *spa* type of CA-MRSA isolates⁹². *Spa* typing is rapid, high throughput with a portable dataset that simplifies information⁷⁶. However, sequencing errors are possible with *spa* typing and it can misclassify particular types due to recombination and/or homoplasy⁷⁷.

Microarrays: Microarrays are widely used to identify single nucleotide polymorphisms (SNPs). There are several microarrays depending on the number and types of DNA spots on the array, the solid support, labeling system, controls for the spots, hybridization conditions, detection system, and data analysis methods. Microarrays (Sam-62) based on 62 *S. aureus* whole genome sequencing (WGS) projects and 153 plasmid sequences shows that MRSA transmission events that are unrecognized by other approaches can be identified and is capable of distinguishing extremely similar but non-identical sequences⁸⁷. Microarray data are reproducible, take 1-2 days and highly accurate. But it is technically demanding, particularly to interpret the data and they do not allow the identification of sequences which are not included in the array⁷⁷.

Whole genome sequencing (WGS): The cost reduction in bench top sequencers and new data analysis softwares makes bacterial WGS possible even in small research and clinical laboratories and it is likely to replace currently used typing methodologies for epidemiological investigations due to its ultimate resolution⁸⁷. WGS of MRSA isolates in an outbreak in neonatal intensive care unit (NICU), Cambridge University Hospitals showed distinct cluster of outbreak isolates (ST22) with a clear separation between these and the non-outbreak isolates based on SNP analysis⁹³. Comparison of WGS of MW2 (USA400) strain and the HA-MRSA strains, Mu50 and N315 revealed that MW2 and the HA-MRSA strains are relatively distant. MW2 strain carry SCCmecIVa element that encodes methicillin resistance gene and penicillinase gene while N315 and Mu30 carries SCCmecII element that had multiple antibiotic resistant genes⁹⁴. WGS has the potential to compare different genomes with a single-nucleotide resolution. This would allow an accurate characterization of transmission events and outbreaks. However, translating this potential into routine practice will involve extensive investigations⁸⁷.

Control measures: When CA-MRSA strains are detected, it is required to do contact tracing to identify and decolonize MRSA carriers among family and close social contacts to limit the community spread of MRSA strains. Screening for MRSA carriage is generally performed by swabbing the anterior nares which has a sensitivity of over 80% that can be increased to 95% when additional sites such as throat, sputum, perineum, and rectum are screened. Decolonization therapy with mupirocin combined with chlorhexidine body washes and oral rifampicin and doxycycline for 5 to 14 days (median 7 days) gives successful MRSA eradication⁶⁶.

CONCLUSION

Rapid methods for MRSA identification and typing are essential to determine the transmission in the community. Molecular typing methods such as SCCmec typing, *spa* typing, MLST and PFGE are recognized as promising methods for MRSA surveillance because of their capability to characterize bacterial chromosomes and mobile genetic elements. Combination of these molecular typing methods is necessary as each method has its own advantages with respect to discriminatory power, rapidity, cost effectiveness, reproducibility, and easy performance.

ACKNOWLEDGEMENT

The authors thank Dr. Kenneth Francis Rodrigues, University Malaysia Sabah for his suggestions.

REFERENCES

- Lowy FD. *Staphylococcus aureus* infections. *N Engl J Med* 1998;**339**(8):520-32.
- Barber M, Rozwadowska-Dowzenko M. Infection by penicillin-resistant staphylococci. *Lancet* 1948;**2**(6530):641-44.
- Panlilio AL, Culver DH, Gaynes RP, *et al.* Methicillin-resistant *Staphylococcus aureus* in US hospitals, 1975-1991. *Infect Control Hosp Epidemiol* 1992;**13**(10):582-86.
- Rosenthal VD, Bijie H, Maki DG, *et al.* International Nosocomial Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004-2009. *Am J Infect Control* 2012;**40**(5):396-07.
- Sowash MG, Uhlemann AC. Community-associated methicillin-resistant *Staphylococcus aureus* case studies. *Methods Mol Biol* 2014; **1085**:25-69.
- Herold BC, Immergluck LC, Maranan MC, *et al.* Community-acquired methicillin-resistant *Staphylococcus aureus* in children with no identified predisposing risk. *JAMA* 1998; **279**:593-98.
- O'Brien FG, Lim TT, Chong FN, *et al.* Diversity among community isolates of methicillin-resistant *Staphylococcus aureus* in Australia. *J Clin Microbiol* 2004;**42**(7):3185-90.
- Millar BC, Loughrey A, Elborn JS, Moore JE. Proposed definitions of community-associated methicillin-resistant *Staphylococcus aureus* (CA-MRSA). *J Hosp Infect* 2007; **67**(2):109-13.
- Udo EE, Pearman JW, Grubb WB. Genetic analysis of community isolates of methicillin-resistant *Staphylococcus aureus* in Western Australia. *J Hosp Infect* 1993;**25**(2):97-108.
- Centers for Disease Control and Prevention (CDC). Four pediatric deaths from community-acquired methicillin-resistant *Staphylococcus aureus*—Minnesota and North Dakota, 1997-1999. *MMWR Morb Mortal Wkly Rep* 1999;**48**(32):707-10.
- Baggett HC, Hennessy TW, Rudolph K, *et al.* Community-onset methicillin-resistant *Staphylococcus aureus* associated with antibiotic use and the cytotoxin Panton-Valentine leukocidin during a furunculosis outbreak in rural Alaska. *J Infect Dis* 2004;**189**(9):1565-73.
- Centers for Disease Control and Prevention (CDC). Community-associated methicillin-resistant *Staphylococcus aureus* infections in Pacific Islanders-Hawaii, 2001-2003. *MMWR Morb Mortal Wkly Rep* 2004;**53**(33):767-70.
- Song JH, Hsueh PR, Chung DR, *et al.* Spread of methicillin-resistant *Staphylococcus aureus* between the community and the hospitals in Asian countries: an ANSORP study. *J Antimicrob Chemother* 2011;**66**(5):1061-69.
- Huang YC, Ho CF, Chen CJ, Su LH, Lin TY. Comparative molecular analysis of community-associated and healthcare-associated methicillin-resistant *Staphylococcus aureus* isolates from children in northern Taiwan. *Clin Microbiol Infect* 2008;**14**(12):1167-72.
- Sam IC, Kahar-Bador M, Chan YF, Loong SK, Ghazali FMN. Multisensitive community-acquired methicillin-resistant *Staphylococcus aureus* infections in Malaysia. *Diagn Microbiol Infect Dis* 2008;**62**(4):437-39.
- Rashid ZZ, Bahari N, Othman A, *et al.* Community-acquired methicillin-resistant *Staphylococcus aureus* in a Malaysian tertiary centre. *Southeast Asian J Trop Med Public Health* 2013;**44**(1):104-08.

17. Ghaznavi-Rad E, Shamsudin MN, Sekawi Z, *et al.* Predominance and emergence of clones of hospital-acquired methicillin-resistant *Staphylococcus aureus* in Malaysia. *J Clin Microbiol* 2010;**48**(3):867-72.
18. Ahmad N, Ruzan IN, Ghani MKA, *et al.* Characteristics of community and hospital acquired methicillin-resistant *Staphylococcus aureus* strains carrying SCCmec type IV isolated in Malaysia. *J Med Microbiol* 2009;**58**(9):1213-18.
19. Chen CJ, Hsueh PR, Su LH, *et al.* Change in the molecular epidemiology of methicillin-resistant *Staphylococcus aureus* bloodstream infections in Taiwan. *Diagn Microbiol Infectious Dis* 2009;**65**(2):199-01.
20. Pannaraj PS, Hulten KG, Gonzalez BE, Mason EO, Kaplan SL. Infective pyomyositis and myositis in children in the era of community-acquired, methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis* 2006;**43**(8):953-60.
21. Adem PV, Montgomery CP, Husain AN, *et al.* *Staphylococcus aureus* sepsis and the Waterhouse-Friderichsen syndrome in children. *N Engl J Med* 2005;**353**(12):1245-51.
22. Miller LG, Perdreau-Remington F, Rieg G, *et al.* Necrotizing fasciitis caused by community-associated methicillin-resistant *Staphylococcus aureus* in Los Angeles. *N Engl J Med* 2005;**352**(14):1445-53.
23. Kravitz GR, Dries DJ, Peterson ML, Schlievert PM. Purpura fulminans due to *Staphylococcus aureus*. *Clin Infect Dis* 2005;**40**(7):941-47.
24. Shopsis B, Kreiswirth BN. Molecular epidemiology of methicillin-resistant *Staphylococcus aureus*. *Emerg Infect Dis* 2001;**7**(2):323-26.
25. Vandenesch F, Naimi T, Enright MC, *et al.* Community-acquired methicillin-resistant *Staphylococcus aureus* carrying Panton-Valentine leukocidin genes: worldwide emergence. *Emerg Infect Dis* 2003;**9**(8):978-84.
26. Boyle-Vavra S, Daum RS. Community-acquired methicillin-resistant *Staphylococcus aureus*: the role of Panton-Valentine leukocidin. *Lab Invest* 2007;**87**(1):3-9.
27. Katayama Y, Ito T, Hiramatsu K. A new class of genetic element, staphylococcus cassette chromosome mec, encodes methicillin resistance in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 2000;**44**(6):1549-55.
28. Katayama Y, Ito T, Hiramatsu K. Genetic organization of the chromosome region surrounding *mecA* in clinical staphylococcal strains: role of IS431-mediated *mecI* deletion in expression of resistance in *mecA*-carrying, low-level methicillin-resistant *Staphylococcus haemolyticus*. *Antimicrob Agents Chemother* 2001;**45**(7):1955-63.
29. IWG-SCC. International Working Group on the Staphylococcal Cassette Chromosome elements [cited 2016 May 17]. Available from: www.SCCmec.org/Pages/SCC_TypesEN.html
30. Ito T, Okuma K, Ma XX, Yuzawa H, Hiramatsu K. Insights on antibiotic resistance of *Staphylococcus aureus* from its whole genome: genomic island SCC. *Drug Resist Updat* 2003;**6**(1):41-52.
31. Lina G, Piémont Y, Godail-Gamot F, *et al.* Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis* 1999;**29**(5):1128-32.
32. Gillet Y, Issartel B, Vanhems P, *et al.* Association between *Staphylococcus aureus* strains carrying gene for Panton-Valentine leukocidin and highly lethal necrotising pneumonia in young immunocompetent patients. *Lancet* 2002;**359**(9308):753-59.
33. Gonzalez BE, Martinez-Aguilar G, Hulten KG, *et al.* Severe staphylococcal sepsis in adolescents in the era of community-acquired methicillin-resistant *Staphylococcus aureus*. *Pediatrics* 2005;**115**(3):642-48.
34. Bocchini CE, Hulten KG, Mason EO, *et al.* Panton-Valentine leukocidin genes are associated with enhanced inflammatory response and local disease in acute hematogenous *Staphylococcus aureus* osteomyelitis in children. *Pediatrics* 2006;**117**(2):433-40.

35. Gerald WT. Medical Importance of the Normal Microform. 1st ed. US: Springer;1999.
36. Williams RE. Skin and nose carriage of bacteriophage types of *Staph aureus*. *J Pathol Bacteriol* 1946;**58**:259-68.
37. Martin TD, Whitehead JE. Carriage of penicillin-resistant *Staph. pyogenes* in healthy adults. *Br Med J* 1949;**1**(4595):173-75.
38. Guinan ME, Dan BB, Guidotti RJ, *et al*. Vaginal colonization with *Staphylococcus aureus* in healthy women: a review of four studies. *Ann Intern Med* 1982; **96**:944-47.
39. Ridley M. Perineal carriage of *Staph. aureus*. *Br Med J* 1959;**1**(5117):270-73.
40. Jacobs SI, Williamson GM, Willis AT. Nasal abnormality and the carrier rate of *Staphylococcus aureus*. *J Clin Pathol* 1961; **14**:519-21.
41. Noble WC, Williams RE, Jevons MP, Shooter RA. Some aspects of nasal carriage of staphylococci. *J Clin Pathol* 1964; **17**:79-83.
42. Kluytmans J, Van Belkum A, Verbrugh H. Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev* 1997; **10**(3):505-20.
43. Lee J, Sung JY, Kim YM, *et al*. Molecular characterization of methicillin-resistant *Staphylococcus aureus* obtained from the anterior nares of healthy Korean children attending daycare centers. *Int J Infect Dis* 2011;**15**(8): e558-63.
44. Ho PL, Chiu SS, Chan MY, *et al*. Molecular epidemiology and nasal carriage of *Staphylococcus aureus* and methicillin-resistant *S. aureus* among young children attending day care centers and kindergartens in Hong Kong. *J Infect* 2012;**64**(5):500-06.
45. Huang YC, Hwang KP, Chen PY, Chen CJ, Lin TY. Prevalence of methicillin-resistant *Staphylococcus aureus* nasal colonization among Taiwanese children in 2005 and 2006. *J Clin Microbiol* 2007;**45**(12):3992-95.
46. Esperson F, Clemmensen I. Isolation of a fibronectin-binding protein from *Staphylococcus aureus*. *Infect Immun* 1982;**37**(2):526-31.
47. Aly R, Shinefield HR, Litz C, Maibach HI. Role of teichoic acid in the binding of *Staphylococcus aureus* to nasal epithelial cells. *J Infect Dis* 1980;**141**(4):463-65.
48. Light IJ, Sutherland JM, Cochran ML, Sutorius J. Ecologic relation between *Staphylococcus aureus* and *Pseudomonas* in a nursery population: Another example of bacterial interference. *N Engl J Med* 1968;**278**(23):1243-47.
49. Aly R, Shinefield HI, Strauss WG, Maibach HI. Bacterial adherence to nasal mucosal cells. *Infect Immun* 1977;**17**(3):546-49.
50. Armstrong-Esther CA, Smith JE. Carriage patterns of *Staphylococcus aureus* in a healthy non-hospital population of adults and children. *Ann Hum Biol* 1976;**3**(3):221-27.
51. Millian SJ, Baldwin JN, Rheins MS, Weiser HH. Studies on the incidence of coagulase-positive staphylococci in a normal unconfined population. *Am J Public Health Nations Health* 1960; **50**:791-98.
52. Kinsman OS, Mckenna R, Noble WC. Association between histocompatibility antigens (HLA) and nasal carriage of *Staphylococcus aureus*. *J Med Microbiol* 1983;**16**(2):215-20.
53. Feinstein V, Musher DM, Cate TR. Bacterial adherence to pharyngeal cells during viral infection. *J Infect Dis* 1980;**141**(2):172-76.
54. Heysell SK, Sheno SV, Catterick K, Thomas TA, Friedland G. Prevalence of methicillin-resistant *Staphylococcus aureus* nasal carriage among hospitalised patients with tuberculosis in rural Kwazulu-Natal. *S Afr Med J* 2011;**101**(5):332-34.
55. Winkler J, Block C, Leibovici L, Faktor J, Pitlik SD. Nasal carriage of *Staphylococcus aureus*: correlation with hormonal status in women. *J Infect Dis* 1990;**162**(6):1400-02.
56. Huijsdens, XW, van Santen-Verheuvcl MG, Spalburg E, *et al*. Multiple cases of familial transmission of community-acquired methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2006;**44**(8):2994-96.

57. De Neeling AJ, Van Leeuwen WJ, Schouls LM, *et al.* Resistance of staphylococci in The Netherlands: surveillance by an electronic network during 1989-1995. *J Antimicrob Chemother* 1998;**41**(1):93-101.
58. Diekema, DJ, Pfaller MA, Schmitz FJ, *et al.* Survey of infections due to *Staphylococcus* species: frequency of occurrence and antimicrobial susceptibility of isolates collected in the United States, Canada, Latin America, Europe, and the Western Pacific region for the SENTRY Antimicrobial Surveillance Program, 1997-1999. *Clin Infect Dis* 2001;**32**(2): S114-32.
59. Urth T, Juul G, Skov R, Schønheyder HC. Spread of a methicillin-resistant *Staphylococcus aureus* ST80-IV clone in a Danish community. *Infect Control Hosp Epidemiol* 2005;**26**(2):144-49.
60. Skov RL, Jensen KS. Community-associated methicillin-resistant *Staphylococcus aureus* as a cause of hospital-acquired infections. *J Hosp Infect* 2009;**73**(4):364-70.
61. Diep BA, Chambers HF, Graber CJ, *et al.* Emergence of multidrug-resistant, community-associated, methicillin-resistant *Staphylococcus aureus* clone USA300 in men who have sex with men. *Ann Intern Med* 2008;**148**(4):249-57.
62. Nguyen DM, Mascola L, Bancroft E. Recurring methicillin-resistant *Staphylococcus aureus* infections in a football team. *Emerg Infect Dis* 2005;**11**(4):526-32.
63. Aiello AE, Lowy FD, Wright LN, Larson EL. Methicillin-resistant *Staphylococcus aureus* among US prisoners and military personnel: review and recommendations for future studies. *Lancet Infect Dis* 2006;**6**(6):335-41.
64. Adcock PM, Pastor P, Medley F, Patterson JE, Murphy TV. Methicillin-resistant *Staphylococcus aureus* in two child care centers. *J Infect Dis* 1998;**178**(2):577-80.
65. Carey AJ, Della-Latta P, Huard R, *et al.* Changes in the molecular epidemiological characteristics of methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit. *Infect Control Hosp Epidemiol* 2010;**31**(6):613-19.
66. Albrich WC, Harbarth S. Health-care workers: source, vector, or victim of MRSA? *Lancet Infect Dis* 2008;**8**(5):289-01.
67. Giuffrè M, Cipolla D, Bonura C, *et al.* Epidemic spread of ST1-MRSA-IVa in a neonatal intensive care unit, Italy. *BMC pediatr* 2012; **12**:64.
68. Kaibni MH, Farraj MA, Adwan K, Essawi TA. Community-acquired methicillin-resistant *Staphylococcus aureus* in Palestine. *J Med Microbiol* 2009;**58**(5):644-47.
69. David MD, Kearns AM, Gossain S, Ganner M, Holmes A. Community-associated methicillin-resistant *Staphylococcus aureus*: nosocomial transmission in a neonatal unit. *J Hosp Infect* 2006;**64**(3):44-50.
70. Al-Tawfiq JA. Father-to-infant transmission of community-acquired methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit. *Infect Control Hosp Epidemiol* 2006;**27**(6):636-37.
71. Ma XX., Sun DD, Wang S, *et al.* Nasal carriage of methicillin-resistant *Staphylococcus aureus* among preclinical medical students: epidemiologic and molecular characteristics of methicillin-resistant *S. aureus* clones. *Diagn Microbiol Infect Dis* 2011;**70**(1):22-30.
72. Chen CS, Chen CY, Huang YC. Nasal carriage rate and molecular epidemiology of methicillin-resistant *Staphylococcus aureus* among medical students at a Taiwanese university. *Int J Infect Dis* 2012;**16**(11): e799-03.
73. Bootsma MCJ, Diekmann O, Bonten MJ. Controlling methicillin-resistant *Staphylococcus aureus*: quantifying the effects of interventions and rapid diagnostic testing. *Proc Natl Acad Sci U S A* 2006;**103**(14):5620-25.
74. Raboud J, Saskin R, Simor A, *et al.* Modeling transmission of methicillin-resistant *Staphylococcus aureus* among patients admitted to a hospital. *Infect Control Hosp Epidemiol* 2005;**26**(7):607-15.
75. Feng Y, Chen CJ, Su LH, *et al.* Evolution and pathogenesis of *Staphylococcus aureus*: lessons learned from genotyping and comparative genomics. *FEMS Microbiol Rev* 2008;**32**(1):23-37.

76. Struelens MJ, Hawkey PM, French GL, Witte W, Tacconelli E. Laboratory tools and strategies for methicillin-resistant *Staphylococcus aureus* screening, surveillance and typing: state of the art and unmet needs. *Clin Microbiol Infect* 2009;**15**(2):112-19.
77. Lindsay JA. Staphylococci. In: de Filippis I, McKee ML, editors. *Molecular Typing in Bacterial Infections*. New York:Springer; 2013. p. 385-405.
78. Malhotra-Kumar S, Haccuria K, Michiels M, *et al.* Current trends in rapid diagnostics for methicillin-resistant *Staphylococcus aureus* and glycopeptide-resistant enterococcus species. *J Clin Microbiol* 2008;**46**(5):577-87.
79. Johnson G, Millar MR, Matthews S, *et al.* Evaluation of BacLite Rapid MRSA, a rapid culture based screening test for the detection of ciprofloxacin and methicillin resistant *S. aureus* (MRSA) from screening swabs. *BMC Microbiol* 2006; **6**:83.
80. McClure JA, Conly JM, Lau V, *et al.* Novel multiplex PCR assay for detection of the staphylococcal virulence marker Pantone-Valentine leukocidin genes and simultaneous discrimination of methicillin-susceptible from -resistant staphylococci. *J Clin Microbiol* 2006; **44**:1141-44.
81. Francois P, Pittet D, Bento M, *et al.* Rapid detection of methicillin-resistant *Staphylococcus aureus* directly from sterile or nonsterile clinical samples by a new molecular assay. *J Clin Microbiol* 2003;**41**(1):254-60.
82. Peterson LR, Petzel RA, Clabots CR, Fasching CE, Gerding DN. Medical technologists using molecular epidemiology as part of the infection control team. *Diagn Microbiol Infect Dis* 1993;**16**(4):303-11.
83. Stefani S, Chung DR, Lindsay JA, *et al.* Methicillin-resistant *Staphylococcus aureus* (MRSA): global epidemiology and harmonisation of typing methods. *Int J Antimicrob Agents* 2012;**39**(4):273-82.
84. Harris SR, Cartwright EJ, Török ME *et al.* Whole-genome sequencing for analysis of an outbreak of methicillin-resistant *Staphylococcus aureus*: a descriptive study. *Lancet Infect Dis* 2013;**13**(2):130-36.
85. Bonofiglio L, Mollerach M, Gardella N. Application of Molecular Typing Methods to the Study of Medically Relevant Gram-Positive Cocci. *Gel Electrophoresis-Advanced Techniques* 2012;113.
86. Mongkolrattanothai K, Boyle S, Kahana MD, Daum RS. Severe *Staphylococcus aureus* infections caused by clonally related community-acquired methicillin-susceptible and methicillin-resistant isolates. *Clin Infect Dis* 2003;**37**(8):1050-58.
87. Sabat AJ, Budimir A, Nashev D, *et al.* Overview of molecular typing methods for outbreak detection and epidemiological surveillance. *Euro Surveill* 2013;**18**(4):20380.
88. Shore AC, Coleman DC. Staphylococcal cassette chromosome mec: recent advances and new insights. *Int J Med Microbiol* 2013;**303**(6-7):350-59.
89. Zhang K, McClure JA, Elsayed S, Louie T, Conly JM. Novel multiplex PCR assay for characterization and concomitant subtyping of staphylococcal cassette chromosome *mec* types I to V in methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 2005;**43**(10):5026-33.
90. Hallin M, Deplano A, Denis O, *et al.* Validation of pulsed-field gel electrophoresis and *spa* typing for long-term, nationwide epidemiological surveillance studies of *Staphylococcus aureus* infections. *J Clin Microbiol* 2007;**45**(1):127-33.
91. Strommenger B, Bräulke C, Heuck D, *et al.* *spa* typing of *Staphylococcus aureus* as a frontline tool in epidemiological typing. *J Clin Microbiol* 2008;**46**(2):574-81.
92. Harmsen D, Claus H, Witte W, *et al.* Typing of methicillin-resistant *Staphylococcus aureus* in a university hospital setting by using novel software for *spa* repeat determination and database management. *J Clin Microbiol* 2003;**41**(12):5442-48.
93. Köser CU, Holden MT, Ellington MJ, *et al.* Rapid whole-genome sequencing for investigation of a neonatal MRSA outbreak. *N Engl J Med* 2012;**366**(24):2267-75.
94. Baba T, Takeuchi F, Kuroda M, *et al.* Genome and virulence determinants of high

virulence community-acquired MRSA.
Lancet 2002;**359**(9320):1819-27.